AD-A066 653

NAVY UNDERWATER SOUND LAB NEW LONDON CONN FURTHER COMMENTS ON BIOLOGICAL BACKSCATTER.(U) NOV 64 B F COLE, T G BELL USL-TM-905-094-64

UNCLASSIFIED









END DATE FILMED 5-79 DDC

F/6 17/1

NL

3 CI AD AO 666

FILE COPY

UNCLASSIFIED MO3 Polect This Copy For -Copy No. USL Problem

> No. 1-001-01-00 (S-F001 03 15-10240)

U. S. NAVY UNDERWATER SOUND LABORATORY FORT TRUMBULL, NEW LONDON, CONNECTICUT

FURTHER COMMENTS ON BIOLOGICAL BACKSCATTER

By B. F. Cole and T. G. Bell

USL Technical Memorandum No. 905-094-64

5 November 1964

INTRODUCTION

On 27 May 1964 Dr. R. P. Chapman of the Naval Research Establishment discussed surface and biological backscatter with T. G. Bell. The results of that discussion were presented in reference (a). Dr. Chapman's more recent work (reference (b)) necessitates a reassessment of the effect of biological backscatter on the SQS-26.

DISCUSSION

Dr. Chapman previously pointed out that the deep scattering layer is always highly frequency dependent, peaking at 5.5 kc in the daytime and at 4 to 5 kc at night (reference (a)). His recent results indicate that it is not unlikely for this scattering peak to shift to the 3 - 4 kc region at night. In one case where the peak shifted to 3.5 kc, the day-night variation was 18 db. The typical DSL surface scattering strength per square yard (integrated over the volume of the layer) was - 41 at night and - 59 in the day. Considering the 4 paths involved (reference (a)), the nighttime level would be equivalent to a sea-surface scattering strength of - 35, that produced by a 20-knot wind for a typical SQS-26 search angle of 15 degrees. This is 8 db higher than previously estimated in

DOWNGRADED AT 3-YEAR INTERVALS DECLASSIFIED AFTER 12 YEARS DOD DIR 5200.10

DISTRIBUTION STATEMENT A

Approved for public release; Distribution Unlimited

1979

UNCLASSIFIED

UNCLASSIFIED

USL Tech Memo No. 905-094-64

reference (a) for the nighttime level.

Since the scattering strength peak frequency depends on the depth of the scatterers, which in turn is believed to correlate with the intensity of light, attempts should be made to correlate the nighttime reverberation background levels with nighttime light intensities. The 3.5 kc resonance which produces the unusually high nighttime levels is probably caused by the scattering layer coming to very shallow depths on dark nights.

RECOMMENDATIONS

It is recommended that light intensity measurements be made during any nighttime reverberation measurements made with the SQS-26 to see if the nighttime reverberation levels are actually dependent on the amount of light. This would be especially important during low wind periods, when the biological scatter is likely to be dominant.

erre	Watte Section	4
106	Buff Saution	0
RANGOUR	TION PAR LATTS	9
OSTIFICAT	TON ATTACK	X
011	FILL	
W		
W	TTIGR/AVAILABILITY 69	
W		
DY	TTIGR/AVAILABILITY 69	n.M.
DY	TTIGR/AVAILABILITY 69	
DY	TTIGR/AVAILABILITY 69	n.M.

B. F. Cole
B. F. Cole
J. S. Bellya

REFERENCES

Bell, T. G., "Discussion with R. P. Chapman of NRE on Surface and Biological Backscattering" (U), USL Technical Memorandum No. 905-058-64, 29 September 1964 (Confidential)

(b) Chapman, R. P., and Marshall, J.R., "Low Frequency Reverberation from Deep Scattering Layers in the Western North Atlantic". Paper G-6, 22nd U. S. Navy Symposium of Underwater Acoustics. October 1964

CONFIDENTIAL

UNCLASSIFIED

2.

USL Tech Memo No. 905-094-64

DISTRIBUTION LIST

External

BUSHIPS (Code 1631) (5 copies) BUSHIPS (Code 1620) (3 copies)

Internal

Code 100	Code 920
101	921
105B	921.4
200	930
900	9304
900A	931.1
900B	931.2
900C	932
902	932.1
904.3	934
905 (Routing Copy)	934.1
9058	935.1
905.1	G. Adkins (Code 932)
905.2	Code 904.2
B. F. Cole (Code 905)	(5 copies)
W. F. Wardle (Code 905)	
Code 907	
907.5	
910	